

LETTER TO THE EDITOR

**Boson realization of non-generic  $sl_q(2)$ - $R$  matrices for the Yang-Baxter equation**

Chang-Pu Sun†‡§, Kang Xue‡§, Xu-Feng Liu§ and Mo-Lin Ge§

† CCAST (World Laboratory), PO Box 8730, Beijing, People's Republic of China

‡ Physics Department, Northeast Normal University, Changchun 130024, People's Republic of China

§ Theoretical Physics Division, Nankai Institute of Mathematics, Tianjin 300071, People's Republic of China||

Received 5 December 1990

**Abstract.** By establishing a new Boson realization of quantum universal enveloping algebra  $sl_q(2)$  and its representations in the non-generic case that  $q$  is a root of unity, we systematically construct non-generic  $R$ -matrices of  $sl_q(2)$  through the universal  $R$ -matrix. These new  $R$ -matrices are not covered by the standard  $R$ -matrices constructed in terms of quantum group and the non-standard ones obtained by using the extended Kauffman's diagram technique.

The Yang-Baxter equation plays a crucial role in nonlinear integrable systems in physics [1-3], and its solutions can be constructed in terms of the quantum universal enveloping algebra (QUEA)  $U_q(L)$  of a classical Lie algebra  $L$  [4-6]. Some results and notions, which will be used in this letter, are briefly reviewed as follows.

Let  $\{e_a\}$  be the basis for a certain Borel subalgebra of  $U_q(L)$  and  $\{e^a\}$  be its dual, then for a given representation  $\rho$  of  $U_q(L)$ , the Hopf algebraic structure of  $U_q(L)$  ensures

$$R = \rho \otimes \rho(\mathcal{R}) = \rho \otimes \rho\left(\sum_a e_a \otimes e^a\right) = \sum_a \rho(e_a) \otimes \rho(e^a)$$

to satisfy the Yang-Baxter equation without spectral parameter

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12} \tag{1}$$

where  $\mathcal{R} = \sum_a e_a \otimes e^a$  is called universal  $R$ -matrix and

$$R_{12} = \sum_a \rho(e_a) \otimes \rho(e^a) \otimes I$$

$$R_{13} = \sum_a \rho(e_a) \otimes I \otimes \rho(e^a)$$

$$R_{23} = \sum_a I \otimes \rho(e_a) \otimes \rho(e^a).$$

|| Mailing address.

If  $\rho$  is chosen to be irreducible,  $R$  is called a standard  $R$ -matrix, which can be expressed by  $q$ - $C$ - $G$  coefficients [7]. In the family of  $R$ -matrices, besides the standard  $R$ -matrices there are non-standard ones associated with Lie algebras  $A_n, B_n, C_n, D_n$ . They are systematically constructed by extending Kauffman's diagram technique [8], and its possible relations to QUEA or quantum group have been analysed [9].

In this letter, we will briefly report the construction of a new class of  $R$ -matrices essentially different from the above mentioned two classes of  $R$ -matrices. Further results based on this letter will be published later.

On the  $q$ -Fock space [10-12]  $\mathcal{F}_q$

$$\{F(n) = a^{+n}|0\rangle|N|0\rangle = a|0\rangle = 0 \quad n \in \mathbb{Z}^+ = \{0, 1, 2, \dots\}$$

where boson operators  $a^+, a = a^-$  and  $N$  satisfy the  $q$ -deformed boson commutation relations

$$aa^+ - q^{-1}a^+a = q^N \quad [N, a^\pm] = \pm a^\pm \quad [a^\pm, a^\pm] = 0 \quad (2)$$

the operators

$$J_+ = \frac{1}{[2]_q} a^{+2} \quad J_- = -\frac{1}{[2]_q} a^2 \quad J_3 = N + \frac{1}{2} \quad (3)$$

satisfy

$$[J_+, J_-] = [J_3]_q^2 \quad [J_3, J_\pm] = \pm 2J_\pm \quad (4)$$

where  $[f]_q = (t^f - t^{-f}) / (t - t^{-1})$ . Thus, (3) gives a new realization of the QUEA  $sl_q(2)$ , which is inhomogeneous and an embedding into the  $q$ -deformed boson realization of  $(C_n)_q$  [13].

A natural representation  $\Gamma$  are defined on  $\mathcal{F}_q$  as

$$J_\pm F(n) = \pm [2]^{-1} ([n][n-1])^{\frac{1}{2}(|\mp|)} F(n \pm 2) \quad J_3 F(n) = (n + \frac{1}{2}) F(n). \quad (5)$$

It is easy to observe that  $\mathcal{F}_q$  and  $\Gamma$  can be reduced as

$$\Gamma = \Gamma^+ \oplus \Gamma^- \quad \mathcal{F}_q = \mathcal{F}_q^+ \oplus \mathcal{F}_q^- \\ \mathcal{F}_q^\pm = \{f^\pm(m) = F(2m + \frac{1}{2}(|\mp|)) | m \in \mathbb{Z}^+\}$$

where  $\Gamma^+$  and  $\Gamma^-$  are representations on invariant subspaces  $\mathcal{F}_q^+$  and  $\mathcal{F}_q^-$  respectively. From (5) we can write down the explicit forms of  $\Gamma^+$  and  $\Gamma^-$

$$J_+ f^\pm(m) = [2]^{-1} f^\pm(m+1) \\ J_- f^\pm(m) = -[2]^{-1} [2m + \frac{1}{2}(|\mp|)] [2m + \frac{1}{2}(|\mp|) - 1] f^\pm(m-1) \quad (6) \\ J_3 f^\pm(m) = (2m + \frac{1}{2}(|\mp|) + \frac{1}{2}) f^\pm(m).$$

It can be proved that  $\Gamma^\pm$  are irreducible when  $q$  is generic and indecomposable (reducible but not completely reducible) when  $q$  is a root of unity [14]. For the non-generic case we have  $q^p = 1$  where  $p$  is an integer larger than or equal to 3, so  $[\alpha p] = 0 (\alpha \in \mathbb{Z}^+)$ , therefore there exist  $\Gamma^\pm$ -invariant subspaces

$$V_{\alpha p}^\pm = \left\{ f^\pm(m) \mid m \geq \frac{\alpha p \pm \sigma(\alpha p)}{3} \right\}$$

determined by the extreme vector  $f^\pm(\frac{1}{2}(\alpha p \pm \sigma(\alpha p)))$  satisfying  $J_- f^\pm(\frac{1}{2}(\alpha p \pm \sigma(\alpha p))) = 0$  where  $\sigma(x) = \frac{1}{2}(1 - (-1)^x)$ .

Now one can easily see that on the quotient spaces  $Q_\alpha^\pm(p) = \mathcal{F}_q^l / V_{\alpha p}^\pm$ :

$$|j, M\rangle^\pm = \tilde{f}^\pm(j+M)|j = \frac{1}{4}(\alpha p \pm \sigma(\alpha p) - 2), M = j, j-3, \dots, j\}$$

$\Gamma^\pm$  induces  $\frac{1}{2}(\alpha p \pm \sigma(\alpha p))$ -dimensional representations of  $\mathfrak{sl}_q(2)$

$$J_+|j, M\rangle^\pm = [2]^{-1}|j, M+1\rangle \quad J_+|j, j\rangle = 0$$

$$J_-|j, M\rangle^\pm = -[2]^{-1}[2(j+M) + \frac{1}{2}(|\mp|)] [2(j+M) + \frac{1}{2}(|\mp|) - 1]|j, M-1\rangle^\pm \quad (7)$$

$$J_3|j, M\rangle^\pm = (2(j+M) + 1 \mp \frac{1}{2})|j, M\rangle^\pm.$$

The above representations are completely new and not covered by the standard angular momentum representation with  $q^p = 1$ , and their relations to the indecomposable representations induced by the regular representation [13] are still unknown.

The explicit matrices of the representation on  $Q_2^+(3)$  are .

$$J_+ = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \quad J_- = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix} \quad J_3 = \frac{1}{2} \begin{bmatrix} 9 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 1 \end{bmatrix}. \quad (8)$$

For  $p=5$  and  $\alpha=1$ , we have the representation

$$J_+ = \begin{bmatrix} 0 & [2]^{-1} & 0 \\ 0 & 0 & [2]^{-1} \\ 0 & 0 & 0 \end{bmatrix} \quad J_- = \begin{bmatrix} 0 & 0 & 0 \\ -1 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix} \quad J_3 = \frac{1}{2} \begin{bmatrix} 9 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (9)$$

on the space  $Q_1^+(5)$ .

Through the universal  $\mathfrak{sl}_q(2)$   $R$ -matrix

$$R = q^{J_3 \otimes J_3} \sum_{n=0}^{\infty} \frac{(1-q^{-4})^n}{[n]_q q^{2!}} q^{n(n-1)} (q^{J_3} J_+ \otimes q^{-J_3} J_-)^n \quad (10)$$

the boson realization  $\mathcal{R}_B \in E_{nd}(\mathcal{F}_q \otimes \mathcal{F}_q)$  of the  $R$ -matrix is given by

$$\mathcal{R}_B = q^{(N+\frac{1}{2}) \otimes (N+\frac{1}{2})} \sum_{n=0}^{\infty} \left\{ \frac{q^{-1}-q}{q+q^{-1}} \right\}^n \frac{q^{n(3n-1)}}{[n]_{q^2}!} a^{+2n} q^{nN} \otimes a^{2n} q^{-nN} \quad (11)$$

in terms of the boson realization (3) of  $\mathfrak{sl}_q(2)$ .

Then, on certain quotient spaces  $Q_\alpha^\pm(p) \otimes Q_\alpha^\pm(p)$  of  $\mathcal{F}_q \otimes \mathcal{F}_q$  we can obtain four classes of  $R$ -matrices from four basic representations on  $Q_\alpha^+(p)$  with even  $\alpha p$ ,  $Q_\alpha^+(p)$  with odd  $\alpha p$ ,  $Q_\alpha^-(p)$  with even  $\alpha p$  and  $Q_\alpha^-(p)$  with odd  $\alpha p$ . For lack of space in this letter, we only write one of them on  $Q_\alpha^+(p)$  with even  $\alpha p$  as follows

$$\begin{aligned} (R)_{m_1 m_2}^{m_1' m_2'} &= q^{(2(m_1+j)+\frac{1}{2})(2(m_2+j)+\frac{1}{2})} \delta_{m_1'}^{m_1} \delta_{m_2'}^{m_2} + \sum_{n=1}^{2j} \frac{(q^{-4}-1)^n}{[2]_q^{2n} [n]_{q^2}!} \\ &\times q^{(2(m_1'+j)+\frac{1}{2})(2(m_2'+j)+\frac{1}{2})+2\Sigma_{i=1}^n (m_1-m_2+2i)} \prod_{l=1}^n [2(m_2+j-l+1)]_q \\ &\times [2(m_2+j-l)+1]_q \delta_{m_1'+n}^{m_1} \delta_{m_2'-n}^{m_2} \end{aligned} \quad (12)$$

It is worth pointing out that only when  $q^p = 1$  will the  $R$ -matrix given by (12) satisfy YBE, so we call it non-generic  $R$ -matrix. In the case of representation (8), (12) gives

a completely new  $R$ -matrix

$$R = q^{5/4} \cdot \left[ \begin{array}{ccccccc} q & & & & & & \\ & q & 0 & & & & \\ & 0 & q & & & & \\ & & & q & q^{-1} - q & 0 & \\ & & & 0 & q^{-1} & 0 & \\ & & & 0 & 0 & q & \\ & & & & & & 1 & q^{-1} - q \\ & & & & & & 0 & 1 \\ & & & & & & & & q^{-1} \end{array} \right] \quad \text{for } j = 1. \quad (13)$$

Of course, for some cases (12) also gives reduced  $R$ -matrices i.e. they can be obtained from standard  $R$ -matrices by letting  $q^p = 1$ . For example, through (12) the representation (9) gives a reduction of the standard  $R$ -matrix with spin 1.

Finally we point out that this letter is only a brief report of the systematic research on non-generic  $R$ -matrices for  $\forall BE$ . The following results are about to be published.

- (1) The general structure of  $R$ -matrix for the indecomposable representation of  $U_q(A_n)$
- (2) The classification of non-generic  $R$ -matrices
- (3) The Yang-Baxterization of non-generic  $R$ -matrices

This work is supported in part by National Foundation of Nature Science of China.

## References

- [1] Yang C N 1967 *Phys. Rev. Lett.* **19** 1312
- [2] Baxter R J 1982 *Exactly Solved Models in Statistical Mechanics* (New York: Academic)
- [3] Yang C N and Ge M L 1989 (eds) *Braid Group, Knot Theory and Statistical Mechanics* (Singapore: World Scientific)
- [4] Drinfeld V G 1986 *Proc. IMC* (Berkeley, CA: University of California Press) p 789
- [5] Jimbo M 1985 *Lett. Math. Phys.* **10**; 1986 *Lett. Math. Phys.* **63** 247; 1986 *Commun. Math. Phys.* **102** 537
- [6] Takhtajan L A and Smirnov F 1990 *Lectures on Quantum Group and Integrable Quantum Field Theory* ed M L Ge and B H Zhao (Singapore: World Scientific)
- [7] Reshetikhin N Yu 1987 LOMI *Preprint* E-4 and E-11
- [8] Kauffman L H 1988 *Ann. Math. Stud.* **115** 1  
Ge M L, Wang L Y, Xue K and Wu Y S 1989 *Int. J. Mod. Phys. A* **4** 3351  
Ge M L, Wang L Y, Li Y Q and Xue K 1990 *J. Phys. A: Math. Gen.* **23** 605  
Ge M L, Li Y Q and Xue K 1990 *J. Phys. A: Math. Gen.* **23** 619
- [9] Ge M L, Sun C P, Wang L Y and Xue K 1990 *J. Phys. A: Math. Gen.* **23** L645
- [10] Biedenharn L C 1989 *J. Phys. A: Math. Gen.* **22** L873
- [11] Sun C P and Fu H C 1989 *J. Phys. A: Math. Gen.* **22** L983
- [12] Macfarlane A J 1989 *J. Phys. A: Math. Gen.* **22** 4551
- [13] Sun C P and Ge M L 1990 *J. Math. Phys.* **31** in press
- [14] Sun C P, Lu J F and Ge M L 1990 *J. Phys. A: Math. Gen.* **23** L1199